

there are faint lines reaching from the A. M. E. down upon the mandibles, sternum reddish brown on the sides, paler in the center; legs reddish, more yellow at tips, a few dark spots at the base of hairs, dorsum of abdomen reddish brown, often with a light median stripe enclosing a spear-mark at base, and a short light mark on each side near tip; venter yellowish with a broad median brown stripe, spinners brown. The ♂ is darker than the ♀. Sometimes the legs are a little banded. Cephalothorax highest at eye-region, gradually sloping concavely until near the posterior end, where it suddenly curves down; legs spiny, abdomen widest in front, tapering to apex; the epigynum consists of a rounded finger, somewhat similar to *O. cinerea* but more slender. Male palpal organ black, the tarsus is more slender than in the other species, the tibia has on the inner side a short pointed projection similar to *O. salticus*, the basal part of the palpal organ is more complicated than in that species.

The four species of *Oxyopes* may be separated by the following table:

1	{ Femora with a black line on under-side	salticus
	{ Femora without a black line on under-side	2
2	{ Abdomen light, with black side and median stripes	scalaris
	{ Abdomen dark, with a few light spots	3
3	{ Dorsum of cephalothorax straight, about as high in middle as in eye-region	cinerea
	{ Dorsum of cephalothorax concave, higher in eye-region than in middle	rufipes



LOCAL ENTOMOLOGICAL NOTES.

Members of the New York Entomological Society and all others, are solicited to contribute to this column, their rare captures, local lists and other items of interest relating to the insect fauna of New York city and vicinity.

LIST OF THE COLEOPTERA OF NORTH EASTERN AMERICA,

WITH SPECIAL REFERENCE TO THE FAUNA OF NEW YORK CITY AND VICINITY.

By CHARLES W. LENG AND WM. BEUTENMULLER.

(CONTINUED FROM PAGE 96.)

SCHIZOGENIUS Putz.

S. planulatus Lee.—N. Y. Taken by Mr. Linell at Coney Island.

S. lineolatus Say.—N. E. Amer. Lives under stones along river banks, June and Sept. Not common in this vicinity.

S. ferrugineus Putz.—N. E. Amer. Occurs in salt marshes. Not common in this vicinity.

S. amphibius Hald.—N. Y., Mo. Taken in this vicinity by Mr. Wm. Jülich.

ARDISTOMIS Putz.

A. obliquata Putz.—N. J. and southward. Taken at Atlantic City, N. J., by Mr. Liebeck.

A. morio Dej.—N. Y. to Fla. Taken in this vicinity by Mr. Jülich.

A. viridis Say.—N. E. Amer. Found in sandy banks. Not common in this vicinity.

A. puncticollis Putz.—Ohio (Dury).

PANAGÆUS Lat.

P. crucigerus Say.—N. Y., N. J. and Southward. Rare in this vicinity. May and June under stones. Taken in April in deserted ant hills (W. B.).

P. fasciatus Say.—N. Y., N. J. and southward. Very rare in this vicinity. May and June.

NOMIUS Lap.

N. pygmæus Dej.—Taken at Newark, N. J., by Mr. Bischoff. Also found in Canada (Harrington).

BEMBIDIUM Lat.

B. punctatostriatum Say.—N. Y., N. J., O. Found along the banks of the Passaic, N. J., in May and Sept.

B. impressum Fabr.—Buffalo, N. Y. *

B. paludosum Sturm.—Buffalo, N. Y.

B. inæquale Say.—N. E. Amer. Found along river banks in May and Sept.

B. nitidulum Dej.—N. E. Amer. Found along the banks of the Passaic, N. J., in May and Sept.

B. nitidum Kby.—Buffalo, N. Y.

B. americanum Dej.—N. E. Amer. Rather common in this vicinity, along banks of ponds. May and Sept.

B. antiquum Dej.—N. Y., N. J. Along river banks, not common in this vicinity.

B. chalceum Dej.—N. Y., N. J., Ohio. Found along the banks of the Passaic in May and Sept.

B. concolor Kby.—Buffalo, N. Y.

B. nigrum Say.—N. Y., N. J., Can. Found along the banks of the Passaic, N. J., in May and Sept.

B. planatum Lee.—N. J. Taken in the same locality as the preceding species.

B. Kuprianovi Mann.—Buffalo, N. Y.

B. fugax Lee.—Buffalo, N. Y., Mo.

B. planum Hald.—Canada, Buffalo, N. Y.

B. rupestre Dej.—N. E. Amer. Common in this vicinity.

B. plagiatum Zimm.—Buffalo, N. Y.

- B. lacunarium** Zimm.—N. Y., N. J. Found along brooks at Fort Lee, N. J. (Linell) and at Coney Island (Leng).
- B. dorsale** Say.—Can., N. Y., N. J., Ohio.
- B. patrulee** Dej.—N. Y., N. J., Ohio, Pa. Found along river banks, May and Sept.
- B. variegatum** Say.—N. E. Amer. Common in this vicinity.
var. posticum Lee.—Taken at Coney Island, N. Y. (Leng).
- B. intermedium** Kby.—N. Y. (Buffalo) N. J., Ohio, Canada.
- B. nigripes** Kby.—Pa., (also Or. Br. Col).
- B. versicolor** Lee.—N. Y., N. J., Ohio. Found along the banks of the Passaic, N. J., in May and Sept.
- B. constrictum** Lee.—N. Y., N. J. Taken at Coney Island (Leng). Common in N. J. Sept.
- B. contractum** Say.—N. Y., N. J. Found along the seashore.
- B. affine** Say.—N. Y., N. J., Ohio.
- B. assimile** Gyll.—Buffalo, N. Y.
- B. quadrimaculatum** Linn.—N. E. Amer. Common everywhere, in fields and gardens.
- B. semistriatum** Hald.—N. Y., N. J. May and Sept.
- B. picipes** Kby.—Ohio (Dury), Peekskill, N. Y. (Sherman), N. Hampshire.
- B. cordatum** Lee.—Ohio (Dury), also Mo.
- B. laevigatum** Lee.—Ohio, (Dury), also Mo.
- B. bifossulatum** Lee.—Peekskill, N. Y., (Sherman).
- B. bimaculatum** Kby.—Ottawa, Canada (Harrington).
- B. pedicellatum** Lee.—Pa. (also Missouri).

TACHYS Schaum.

- T. proximus** Say.—Ohio, Buffalo, N. Y., N. J., Pa. Found along the banks of the Passaic, N. J. May and Sept.
- T. scitulus** Lee.—Ohio, Buffalo, N. Y., N. J. Found in the same locality as the preceding.
- T. occultator** Casey.—N. J. Taken at Brigantine Beach by Dr. Hamilton in September.
- T. laevus** Say.—Ohio, Buffalo, N. Y., N. J., Pa. Rare in this vicinity.
- T. nanus** Gyll.—N. E. Amer. Common in this vicinity under bark of trees; runs very fast.
- T. flavicauda** Say.—N. E. Amer. Common under bark of trees. May and Septembér.
- T. tripunctatus** Say.—N. Y., N. J., Pa. Taken along the banks of the Passaic, N. J., in May and Sept.
- T. vivax** Lee.—Ohio, Buffalo, N. Y., Pa.
- T. capax** Lee.—N. J. and westward. Found in cranberry bogs in N. J. (Smith).
- T. xanthopus** Say.—N. Y., N. J. and westward.
- T. incurvus** Say.—N. E. Amer. Common in the vicinity of N. Y. City.
- T. fuscicornis** Chd.—N. J.
- T. pulchellus** Lee.—Ohio and westward.
- T. ventricosus** Lee.—Peekskill, N. Y. (Sherman).

PATROBUS Dej.

P. longicornis Say.—N. E. Amer. Found in woods in moist places.

P. septentrionis Dej.—Northern States and Europe. Not found in this vicinity.

P. rugicollis Rand.—N. Y., Pa., Mass. and northward. Not found in this vicinity.

PERICOMPSUS Lec.

P. ephippiatus Say.—N. Y. and westward. Found in sandy places near the Hudson River, N. Y. (Sherman).

POGONUS Dej.

P. texanus Chd.—N. J. Rare at Atlantic City.

TRECHUS Clair.

T. rubens Fabr.—Nova Scotia, N. Europe.

T. chalybaeus Mann.—Alaska to N. Hampshire.

MYAS Dej.

M. cyanescens Dej.—N. Y., N. J. Taken in this vicinity at Fort Lee and Clifton, N. J., and on Staten Island, in June and July. Rare.

M. coracinus Say.—N. Y., N. J., Ohio. Rare.

PTEROSTICHUS Bon.

P. adoxus Say.—N. E. Amer. Not common in this vicinity.

P. rostratus Newm.—Can., N. Y. and southward.

P. diligendus Chaud.—N. E. Amer. Not common in this vicinity.

P. honestus Say.—N. E. Amer. Not rare in this vicinity.

P. lachrymosus Newm.—N. Y. and southward. Not rare in this vicinity.

P. stygicus Say.—N. E. Amer. Abundant in this vicinity.

P. coracinus Newm.—N. E. Amer. Not rare at Snake Hill, and Short Hills, N. J.

P. relictus Newm.—N. Y., Pa., Mich. Rare.

P. superciliosus Say.—N. Y., Pa., Mich. Rare.

P. moestus Say.—N. Y., N. J., Pa., W. Va. Rare in this vicinity.

P. punctatissimus Rand.—Mass. and Westward.

P. sculptus Lec.—N. Y. and southward. Taken at Atlantic City, N. J. (Smith).

P. Hamiltoni Horn.—Pa., E. Ohio, Md., W. Va. Not common.

P. Sayi Brulle.—N. E. Amer. Common in this vicinity.

P. lucublandus Say.—N. E. Amer. Very common everywhere.

P. ebeninus Dej.—N. Y. and southward. Rare in this vicinity.

P. caudicalis Say.—N. E. Amer. Rare in this vicinity.

P. luctuosus Dej.—N. E. Amer. Taken at Snake Hill, N. J. Rare.

P. corvinus Dej.—N. E. Amer. Not common in this vicinity.

P. purpuratus Lec.—N. Y. and westward. Not found in this vicinity. Rare.

- P. mutus *Say*.—N. E. Amer. Very common everywhere in this vicinity.
P. pennsylvanicus *Lec.*.—N. E. Amer. Not found in this vicinity.
P. orinomum *Leach*.—Canada, Nova Scotia.
P. Luczotii *Dej.*.—N. Y. and northward. Not found in this vicinity.
P. erythropus *Dej.*.—N. E. Amer. Rare in this vicinity.
P. patruelis *Dej.*.—N. E. Amer. Common in this vicinity.
P. femoralis *Kby.*.—Can., N. V. and westward. Wis. (O. Dietz).
P. corrusculus *Lec.*.—N. Y., Mass. Rare under moss.
P. mandibularis *Kby.*.—N. Hampshire, Can.
P. Haldemani *Lec.*.—N. Y. and southward. Rare on Staten Island (Leng).
P. tartaricus *Say*.—N. Y., southward and westward. Taken at Hoboken,
N. J. and Astoria, L. I. May.
P. scrutator *Lec.*.—N. V., Can., Ill.
P. gravis, *Lec.*.—Pa.
P. permundus *Say*.—Ohio.
P. substriatus *Lec.*.—N. J. (Davis), N. Y. (Jülich) and southward.
P. rotundatus *Lec.*.—N. Y. (Jülich), Ga. (Lec).
P. unicolor *Say*.—N. Y. and southward. Not found in this vicinity.
P. approximatus *Lec.*.—Pa. and southward.
P. hudsonicus *Lec.*.—Labrador.

EVARTHRUS *Lec.*

- E. sigillatus *Say*.—N. J. and southward.
E. sodalis *Lec.*.—Pa. and southward. Wis., Ohio (Dietz).
E. furtivus *Lec.*.—Pa. and southward.

AMARA Bon.

- A. avida *Say*.—N. E. Amer. Rare in this vicinity.
A. arenaria *Lec.*.—Buffalo, N. Y., N. Hampshire.
A. exarata *Dej.*.—N. E. Amer. Not common.
A. latior *Kby.*.—N. J.
A. angustata *Say*.—N. E. Amer. Not rare in this vicinity.
A. pallipes *Kby.*.—N. J. and westward.
A. impuncticollis *Say*.—N. E. Amer. Common in this vicinity.
A. basillaris *Say*.—N. Y., N. J. (Linell).
A. cupreolata *Putz.*.—N. J., Ill.
A. interstitialis *Dej.*.—Can., N. V., N. J. Not rare in this vicinity.
A. obesa *Say*.—N. Y., N. J., Can. and westward.
A. fallax *Lec.*.—Buffalo, N. Y. and westward.
A. erratica *Sturm*.—Can., Vt., Mich.
A. chalcea *Dej.*.—N. V., N. J. and westward.
A. gibba *Lec.*.—N. J.
A. rubrica *Hald.*.—N. V., N. J. Rare in this vicinity.
A. subænea *Lec.*.—Can., N. V., N. J.
A. musculus *Say*.—N. V., N. J. and westward. Not common in this
vicinity.
A. acutangula *Putz.*.—N. J. Rare.
A. similis *Kby.*.—Labrador.

- A. polita** *Lec.*—Can., N. Y. (O. Dietz).
A. septentrionalis *Lec.*—N. Y. (O. Dietz).
A. hyperborea *Dej.*—Labrador.

LOXANDRUS *Lec.*

- L. agilis** *Dej.*—Pa. and southward.

DIPLOCHILA *Brulle.*

- D. laticollis** *Lec.*—New York and westward. Not rare in moist places, in this vicinity. May and June.
var. major *Lec.*—Found with the preceding.
D. impressicollis *Dej.*—Can., N. Y. and westward.

DICÆLUS *Bon.*

- D. dilatatus** *Say.*—N. Y. and westward. Rare in this vicinity. May and June.
D. purpuratus *Bon.*—N. Y., Mass., southward and westward.
D. sculptilis *Say.*—Ohio, southward and westward. Wis. (O. Dietz).
D. furvus *Dej.*—Ohio, Southward and westward.
D. crenatus *Lec.*—A southern species on the Buffalo list.
D. ovalis *Lec.*—N. Y., west and southward. Taken at Snake Hill, N. J.
D. elongatus *Bon.*—N. E. Amer. Not rare in this vicinity.
D. ambiguus *Laf.*—N. Y., west and south. Not found in this vicinity.
D. teter *Bon.*—N. E. Amer. Not rare along the Palisades, N. J.
D. politus *Dej.*—N. E. Amer. Not rare in this vicinity.

LICINUS *Lat.*

- L. silphoides** *Fab.*—Mass. and Europe.

BADISTER *Clairv.*

- B. notatus** *Hald.*—Can., N. Y., N. J., westward and southward. Rare in this vicinity.
B. micans *Lec.*—Mass., N. Y., N. J., Ill., Fla. Rare in this vicinity.
B. pulchellus *Lec.*—Can., N. Y., Ill., Ind., Fla.
B. maculatus *Lec.*—Pa.
B. ferrugineus *Dej.*—Cal., Alaska (on the Buffalo list).
B. flavipes *Lec.*—N. Y., La., Fla.
B. reflexus *Lec.*—N. Y., Mich., La.

CALATHUS *Bon.*

- C. gregarius** *Say.*—Can., N. Y. to Fla. to Tex. Common in various localities in this vicinity.
C. ingratus *Dej.*—Can., Labrador, Alaska.
C. opaculus *Lec.*—N. E. Amer. Rare in this vicinity.
C. advena *Lec.*—Alaska to N. Hampshire.
C. impunctatus *Say.*—N. E. Amer. Not rare in this vicinity, especially at the sea shore.

PLATYNUS Bon.

P. angustatus Dej.—Middle and Southern States. Rare at Fort Lee, N. J.

P. caudatus Lec.—Ohio.

P. hypolitus Say.—Middle and Southern States.

P. decens Say.—Can., to N. C. Scarce in this vicinity.

P. sinuatus Dej.—N. E. Amer. Not common in this vicinity.

P. opaculus Lec.—Middle and Western States. Rare at Fort Lee, N. J.

P. tenuicollis Lec.—Lake Superior. Taken at Atlantic City, N. J. (Castle).

P. cincticollis Say.—Middle and Southern States. Common under stones along brooks.

P. reflexus Lec.—Can., and Middle States. Found in same situation as the preceding species.

P. extensicollis Say.—N. E. Amer. Common under stones along brooks and rivers in this vicinity.

var. viridis Lec.—Southern and Western States. (On the Buffalo list).

P. decorus Say.—N. E. Amer. Not rare along the Passaic, N. J.

F. anchomenoides Rand.—Can., and Maine.

P. clemens Lec.—Nova Scotia.

P. errans Say.—Canada.

P. moerens Dej.—Can., and Atlantic States.

P. pusillus Lec.—Can., N. Y. and westward.

P. collaris Say.—Southern States. (On the Buffalo list).

P. tenuis Lec.—Can., and Middle States.

P. atratus Lec.—N. E. Amer. Rare in this vicinity.

P. melanarius Dej.—N. E. Amer. Common in this vicinity.

P. propinquus G. & H.—Can., to Mass., N. J.

P. affinis K'hy. N. E. Amer. Rare in this vicinity.

P. corvus Lec.—Can., to Or.

P. metallescens Lec.—Lake Superior to N. J. Not rare in this vicinity.

P. deceptivus Lec.—Nova Scotia to Lake Superior.

P. cupripennis Say.—N. E. Amer. Common in this vicinity.

var. nitidulum Dej.—Bear Lake, Pa.

P. Hardyi Lec.—Newfoundland.

P. excavatus Dej.—Can., Middle and Western States. Rare in this vicinity.

P. ferreus Hald.—Middle and Western States. Not rare along the Palisades, N. J.

P. basalis Lec.—Pa., N. J., to Fla. Very rare. Taken at Hoboken, N. J. (Linell).

P. maculicollis Dej.—Cal. (On the Buffalo list).

P. variolatus Lec.—Cal. (On the Buffalo list).

P. nutans Say.—Can., N. Y. to Fla. Rare in this vicinity.

P. octopunctatus Fabr.—N. E. Amer. Rare in this vicinity.

P. placidus Say.—N. E. Amer. Common.

P. vicinus G. & H.—Can., Lake Superior.

P. obsoletus Say.—N. E. Amer. Rather common in this vicinity.

P. quadripunctatus Dej.—Can., Alaska, N. Y. (On the Buffalo list).

P. æruginosus Dej.—Atlantic States. Not common under bark in this vicinity.

P. crenistriatus *Lee.*—N. Y. and westward. Abundant on seashore of Long Island and New Jersey.

P. rubripes *Zimm.*—Middle and Western States. Not common in this vicinity.

P. punctiformis *Say.*—N. E. Amer. Rather common in this vicinity.

P. sordens *Kby.*—Can., to N. J. Rare in this vicinity.

P. picicornis *Lee.*—Can., N. Y. and westward. Not found around N. Y. City.

P. ruficornis *Lee.*—N. Y. and Westward. Wis. (O. Dietz).

P. retractus *Lee.*—Can., Pa., Mass. Lake Superior.

P. picipennis *Kby.*—Mass., N. Y., N. J. and northward.

P. lutulentus *Lee.*—N. E. Amer. Not common in this vicinity.

P. exaratus *Mauu.*—Pa.

P. limbatus *Say.*—Can. (Coupe). Mo.

OLISTHOPUS *Dej.*

O. parvatus *Say.*—N. E. Amer.

O. micans *Lee.*—N. J. southward and westward.

ATRANUS *Lee.*

A. pubescens *Dej.*—N. Y. and westward.

LEPTOTRACHELUS *Lat.*

L. dorsalis *Fab.*—N. Y. and westward.

CASNONIA *Lat.*

C. pennsylvanica *L.*—N. E. Amer.

C. ludoviciana *Salle.*—D. C. and South.

GALERITA *Fab.*

G. janus *Fab.*—N. E. Amer. Common everywhere.

G. bicolor *Dru.*—N. Y., N. J., Ga., Mo.

TETRAGONODERUS *Dej.*

T. fasciatus *Hald.*—N. Y. and westward.

LEBIA *Lat.*

L. grandis *Hentz.*—N. E. Amer. Common everywhere.

L. atriventris *Say.*—N. E. Amer. Not as common as the preceding species.

L. tricolor *Say.*—N. E. Amer.

L. pulchella *Dej.*—N. E. Amer. Not rare.

L. cyanipennis *Dej.*—N. J., Cal.

L. marginicollis *Dej.*—N. J. Rare.

L. viridis *Say.*—N. E. Amer. Common.

L. moesta *Lee.*—N. Y., N. J. and westward.

- L.** *pumila* *Lee.*—N. E. Amer. Not common.
L. *pleuritica* *Lee.*—N. E. Amer. Not common.
L. *viridipennis* *Dej.*—N. E. Amer.
L. *lobulata* *Lee.*—Ohio, Va., La.
L. *ornata* *Say.*—N. E. Amer.
L. *collaris* *Dej.*—Middle and Southern States.
L. *analis* *Dej.*—N. E. Amer.
L. *fuscata* *Dej.*—N. E. Amer.
var. frigida *Chaud.*—Mass.
L. *scapularis* *Dej.*—N. E. Amer.
L. *vittata* *Fab.*—N. Y., N. J., Pa. to Tex. Not common in this vicinity.
var. Spraguei Horn.—N. J. (Castle), Tex.
L. *furcata*—*Lee.* N. E. Amer.
L. *bivittata* *Fab.*—Mass., N. Y., N. J. Westward and Southward.

COPTODERA *Dej.*

- C.** *aerata* *Dej.*—N. E. Amer. Rare.

DROMIUS *Bon.*

- D.** *piceus* *Dej.*—Mass., N. Y., Pa. to Cal. Found under bark and on flowers.

APRISTUS *Chd.*

- A.** *cordicollis* *Lee.*—N. Y., N. J.
A. *subsulcatus* *Dej.*—N. Y., Can.

BLECHRUS *Mots.*

- B.** *nigrinus* *Maun.*—Can., N. Y. and westward. Found under bark.

METABLETUS *Sch.-Goch.*

- M.** *americanus* *Dej.*—N. E. Amer.

AXINOPALPUS *Lee.*

- A.** *biplagiatus* *Dej.*—Can., Mass. to Calif.

CALLIDA *Dej.*

- C.** *viridipennis* *Say.*—N. Y. to Texas.
C. *punctata* *Lee.*—N. J., N. Y., Mich., O., La., Ks., Can.
C. *purpurea* *Say.*—N. J., N. Y., Mich., Ga., Mo., Ks., Neb.

PLOCHIONUS *Dej.*

- P.** *Bonfilsii* *Dej.*—Pa., Mex.
P. *timidus* *Hald.*—N. Y., N. J., Ala., Cal. Under bark.
P. *amandus* *Hald.*—N. Y. and Southward.

PINACODERA *Schaum.*

P. limbata *Dej.*—N. Y., N. J.

var. fuscata *Dej.*—N. Y.

P. platicollis *Say.*—N. Y., N. J., Fla., Tex.

CYMINDIS *Lat.*

C. cribicollis *Dej.*—N. Y. and westward.

C. elegans *Lee.*—N. Y.

C. unicolor *Kby.*—L. Sup., Labrador.

C. americana *Dej.*—N. Y., N. J. and westward.

C. pilosa *Lee.*—Can., N. Y., N. J. and westward.

C. borealis *Lee.*—Can., Nova Scotia.

C. neglecta *Hald.*—Can., N. Y., N. J., Pa.

APENES *Lee.*

A. lucidula *Dej.*—N. Y., N. J., Fla. Seashore.

A. sinuata *Say.*—N. Y., N. J.

HELLUOMORPHA *Lat.*

H. bicolor *Harr.*—Mass. to Fla. and westward. Taken at Moshulu, N. Y., and Fort Lee, N. J. Rare.

H. ferruginea *Lee.*—N. Y. to Tex. Rare in this vicinity.

BRACHYNUS *Web.*

B. viridipennis *Dej.*—N. Y., N. J.

B. minutus *Harr.*—N. Y., N. J.

B. perplexus *Dej.*—Can., N. Y., N. J. and westward.

B. medius *Harr.*—N. E. Amer.

B. quadripennis *Dej.*—N. Y., N. J., Pa., Mass.

B. conformis *Dej.*—N. Y., N. J.

B. cyanipennis *Say.*—N. Y., N. J.

B. alternans *Dej.*—Can., N. Y., N. J. and westward.

B. fumans *Fab.*—N. E. Amer.

var. similis *Lee.*—N. E. Amer.

B. cordicollis *Dej.*—Can., N. Y., N. J. and westward.

B. americanus *Lee.*—Can., N. Y. and westward.

B. ballistarius *Lee.*—Can., Ill., Mo.

CHLAENIUS *Bon.*

C. erythropus *Germ.*—Ohio to Neb., La. Rare in N. J., Buffalo, N. Y., Fla. (O. Dietz).

C. fuscicornis *Dej.*—Gulf States to Ill. (On the Buffalo list.)

C. sericeus *Forst.*—N. E. Amer. Common everywhere.

C. laticollis *Say.*—N. Y. to Fla. and westward.

C. diffinis *Chd.*—Middle States Region.

C. aestivus *Say.*—N. E. Amer.

C. prasinus *Dej.*—Middle States to Col.

C. leucoscelis *Chev.*—N. E. Amer. Under stones by Croton River, N. Y.

- C. solitarius** *Say.*—N. E. Amer. Not found in this vicinity.
C. vafer *Lec.*—Texas. (On the Buffalo list.)
C. nemoralis *Say.*—N. E. Amer. Common.
C. tricolor *Dej.*—N. E. Amer. Common everywhere.
C. pennsylvanicus *Say.*—N. Y., N. J. and northward.
C. impunctifrons *Say.*—N. E. Amer. Not common.
C. niger *Rand.*—N. E. Amer. Rare.
C. purpuricollis *Rand.*—N. E. Amer. Very rare.
C. tomentosus *Say.*—N. E. Amer. Common.
C. platyderus *Chd.*—Ill., west and southward.

ANOMOGLOSSUS *Chd.*

- A. emarginatus** *Say.*—N. E. Amer. Not rare in this vicinity.
A. pusillus *Say.*—N. E. Amer. Rare in this vicinity.

BRACHYLOBUS *Chd.*

- B. lithophilus** *Say.*—N. E. Amer. Rare in this vicinity.

LACHNOCREPIS *Lec.*

- L. parallelus** *Say.*—N. Y. and westward.

OODES *Bon.*

- O. amaroides** *Dej.*—N. Y., N. J., Pa., southward and westward. Very rare in this vicinity.

- O. americanus** *Dej.*—N. Y. and southward.
O. fluvialis *Lec.*—Can., N. Y., southward and westward.
O. Lecontei *Chd.*—Ia., (On the N. J. list.)
O. cupraeus *Chd.*—Tex., Mo. (On the Buffalo list.)

GEOPINUS *Lec.*

- G. incrassatus** *Dej.*—N. Y., N. J. and westward. Sandy districts near water, often six inches deep.

NOTHOPUS *Lec.*

- N. zabroides** *Lec.*—Ill. (O. Dietz.)

CRATACANTHUS *Dej.*

- C. dubius** *Dej.*—N. Y., N. J. and westward. Found in sandy places, near water.

AGONODERUS *Dej.*

- A. lineola** *Fab.*—N. E. Amer. Common everywhere.
A. infuscatus *Dej.*—N. Y. and southward.
A. pallipes *Fab.*—N. E. Amer. Common everywhere.
A. partiarius *Say.*—N. E. Amer.
A. pauperculus *Dej.*—Southern States. (On the N. J. and Buffalo lists).
A. indistinctus *Dej.*—N. Y., N. J.
A. testaceus *Dej.*—N. Y., N. J.
A. micros *Lec.*—Eastern States.

DISCODERUS *Lee.*

- D. parallelus** *Hald.*—N. J., southward and westward. Seashore.
D. impotens *Lee.*—Ohio. (O. Dietz.)

GYNANDROPUS *Dej.*

- G. hylacis** *Say.*—N. Y., N. J. Under bark.

HARPALUS *Lat.*

- H. dichrous** *Say.*—N. V., N. J.
H. vulpeculus *Say.*—N. V., N. J.
H. autumnalis *Say.*—N. V., N. J. Rare at seashore.
H. erraticus *Say.*—Can., N. H., Mass., N. Y., N. J.
H. viridianeus *Beaum.*—N. E. Amer.
H. basilaris *Khr.*—Can.
H. laticeps *Lee.*—Wis. (O. Dietz.) Can.
H. caliginosus *Fab.*—N. E. Amer. (U. S.)
H. faunus *Say.*—N. E. Amer.
H. convivus *Lee.*—N. J. Rare.
H. vagans *Lee.*—N. E. Amer.
H. pennsylvanicus *Dej.*—N. E. Amer. [U. S.]
var. **compar** *Lee.*—N. E. Amer.
var. **erythropus** *Dej.*—N. E. Amer.
H. spadiceus *Dej.*—N. J. [rare] and westward.
H. fallax *Lee.*—N. V. [Linell], N. J.
H. pleroticus *Khr.*—N. V., N. J. and westward.
H. herbivagus *Say.*—N. E. Amer.
H. nitidulus *Chd.*—N. V., N. J. Rare at seashore.
H. innocuus *Lee.*—Taken in this vicinity by Mr. Linell.
H. rufimanus *Lee.*—Pa.
H. gravis *Lee.*—Tex. [On the Buffalo list.]

SELENOPHORUS *Dej.*

- S. pedicularius** *Dej.*—N. V. and southward. Seashore in Sept.
S. iripennis *Say.*—Ill., Ga., Tex. [On the Buffalo list.]
S. gagatinus *Dej.*—Mass. and westward.
S. opalinus *Lee.*—N. E. Amer. Seashore, Sept.
S. ovalis *Dej.*—N. J., Ga., Fla. Seashore, Sept.
S. ellipticus *Dej.*—N. V., N. J., Ga., Tex. Seashore, Sept.

STENOLOPHUS *Dej.*

- S. carbonarius** *Brulle.*—Middle and Southern States. Seashore, Sept.
S. spretus *Dej.*—N. V. and southward.
S. fuliginosus *Dej.*—N. V., N. J. and westward.
S. plebeius *Dej.*—N. V., N. J. Seashore, Sept.
S. conjunctus *Say.*—N. E. Amer. [U. S.]
S. ochropeplus *Say.*—N. E. Amer.
S. dissimilis *Dej.*—Can., N. J., southward and westward.
S. alternans *Lee.*—Pa.

ACUPALPUS *Lat.*

- A. *hydropicus* *Lec.*—N. Y., N. J. May.
A. *carus* *Lec.*—Can., N. J., N. Y.

BRADYCELLUS *Er.*

- B. *linearis* *Lec.*—Pa., Wisc.
B. *cognatus* *Gyll.*—Europe and Northern America.
B. *neglectus* *Lec.*—Buffalo and northwest.
B. *rupestris* *Say.*—N. E. Amer.
B. *tantillus* *Dej.*—U. S.
B. *nigriceps* *Lec.*—N. Y., Va.

TACHYCELLUS *Moraw.*

- T. *nigrinus* *Dej.*—Alaska to Can.
T. *Kirbyi* *Horn.*—Canada to Ohio.
T. *atrimedius* *Say.*—N. E. Amer.
T. *badiipennis* *Hald.*—N. E. Amer.

ANISODACTYLUS *Dej.*

- A. *dulcicollis* *Laf.*—N. J. and southward.
A. *rusticus* *Say.*—N. E. Amer. Common everywhere.
A. *carbonarius* *Say.*—N. E. Amer.
A. *interpunctatus* *Kby.*—N. E. Amer.
A. *agricola* *Say.*—N. Y., southward and westward.
A. *Harrisii* *Lec.*—N. E. Amer.
A. *nigerimus* *Dej.*—N. E. Amer.
A. *nigrita* *Dej.*—N. E. Amer.
A. *melanopus* *Hald.*—N. J., Pa., Ill.
A. *discoideus* *Dej.*—N. E. Amer. Rare.
A. *baltimorensis* *Say.*—N. E. Amer. Common.
A. *laetus* *Dej.*—N. Y., N. J., Ga., Tex.
A. *coenus* *Say.*—N. E. Amer.
A. *sericeus* *Harr.*—N. E. Amer.
A. *lugubris* *Dej.*—N. J., Ia., Can., Mo.
A. *interstitialis* *Say.*—N. E. Amer.
A. *verticalis* *Lec.*—N. E. Amer.
A. *terminatus* *Say.*—N. Y., N. J. and westward.
A. *piceus* *Lec.*—N. E. Amer.
A. *semipunctatus* *Lec.*—Or. [On the Buffalo list.]
A. *punctulatus* *Lec.*—Middle State Region.

(TO BE CONTINUED.)